

Conformal Invariance, Anomalous Gravitational Coupling and Derivation of a Particle Concept

Hadi Salehi *

Institute for Studies in Nonlinear Analysis, School of Mathematical Sciences,
Shahid Beheshti University, P.O.Box 19395-4716, Evin, Tehran 19834, Iran.

and

Department of Physics, Shahid Beheshti University, Evin, Tehran 19839, Iran.

Yousef Bisabr[†]

Department of Physics, Shahid Rajaee University, Lavizan, Tehran 16788, Iran.

Abstract

We study a gravitational model whose vacuum sector is invariant under conformal transformations. In this model we investigate the anomalous gravitational coupling of the large-scale matter. In this kind of coupling the large-scale matter is taken to couple to a metric which is different but conformally related to the metric appearing explicitly in the vacuum sector. The effect of the conformal symmetry breaking of the large-scale matter would lead in general to a variable strength of the anomalous gravitational coupling. This feature is used to derive a relativistic particle concept which shares the essential dynamical characteristics of the particle concept used in the causal interpretation of quantum mechanics with respect to the form of the Hamilton-Jacobi equation. The basic aspect of this result is that it relates the variable strength of the anomalous gravitational coupling of the large-scale matter to the appearance of a term similar to a quantum potential term. Some of the general characteristics of the corresponding pilot wave are discussed.

*e-mail: h-salehi@cc.sbu.ac.ir.

[†]e-mail: y-bisabr@srttu.edu.

The principle of conformal invariance requires that all the fundamental equations of physics should be invariant under local (spacetime dependent) changes of units of length and time [1]. This requirement poses a fundamental symmetry in physical theories which is of ever-increasing interest. The establishment of such a symmetry in the vacuum sector of a gravitational model leads one to face with a problem concerning the incorporation of matter to such models. In fact, once the vacuum sector is taken to be conformally invariant, all the conformal frames of the metric should be considered as dynamically equivalent. By implication there is no *a priori* evidence determining to which of these frames the matter couples. From a classical point of view one may appeal to the weak equivalence principle which forces us to consider the coupling of matter to the metric which enters explicitly the action describing the vacuum sector. This is what is generally known as the normal coupling. At the quantum level, however, there is not a direct evidence demonstrating the validity of the weak equivalence principle. Thus questions such as what is the nature of gravitational coupling of matter at this level and what are the physical consequences of this coupling are of obvious theoretical significance.

In this paper we deal with questions of this type in the framework of a conformal invariant gravitational model in which the gravitational interaction of matter can be described both by the normal coupling and the anomalous coupling. The anomalous coupling means that the metric tensor entering the gravitational part and that entering the matter part of the action correspond to different conformal frames. We shall establish such a distinction by introducing a nontrivial conformal factor which in our model is considered as a dynamical field. We show that the emergence of this field can be used to model a relativistic particle concept which shares the essential characteristics of the particle concept in the casual interpretation of quantum mechanics [2] with respect to the form of the Hamilton Jacobi equation. This deduction of the particle concept is carried out on the basis of a conformal symmetry breaking.

It should be noted that the relation between the conformal invariant gravitational models and causal interpretation of quantum mechanics has been studied in many works, see, e.g [3] [4] [5]. The present work deals with the exploration of the role played by the anomalous gravitational coupling in this context. Throughout the following we shall use units in which $\hbar = c = 1$ and the signature is $(-+++)$.

Consider the vacuum sector of a scalar tensor theory defined by the conformal invariant gravitational action functional

$$S = \frac{1}{2} \int d^4x \sqrt{-g} (g^{\mu\nu} \nabla_\mu \phi \nabla_\nu \phi + \frac{1}{6} R \phi^2) , \quad (1)$$

where ϕ is a scalar field, R is the curvature scalar associated with the metric tensor $g_{\mu\nu}$ and ∇_μ denotes a covariant differentiation.

The conformal invariance of the action (1) means that it is invariant under conformal transformations

$$g_{\mu\nu} \rightarrow \Omega^2 g_{\mu\nu} , \quad (2)$$

$$\phi(x) \rightarrow \Omega^{-1} \phi(x) , \quad (3)$$

where Ω is a smooth dimensionless spacetime function. The implication is that in the vacuum sector of a gravitational model defined by the action (1) the conformal frame of the metric remains dynamically unspecified. This point is very important concerning the gravitational

coupling of a matter system. In fact if one wants to add a matter action to the gravitational action (1) one has to decide which conformal frame of the metric is chosen for the description of the matter action. The dynamical characteristics of a matter system may depend significantly on the choice of this conformal frame.

To incorporate a matter system one may assume that the conformal frame of the metric used in the matter action is the same as that of the gravitational action (1). This corresponds to the normal gravitational coupling and reflects one of the main characteristics in the standard description of matter systems. For our purpose we need to consider besides the normal coupling the anomalous coupling in which case one distinguishes between the conformal frame of the metric used in the gravitational action (1) and that used in the matter action. In explicit terms we intend to investigate a biscalar-tensor theory defined by the action functional

$$S' = \frac{1}{2} \int d^4x \sqrt{-g} \{ g^{\mu\nu} \nabla_\mu \phi \nabla_\nu \phi + (g^{\mu\nu} \nabla_\mu \sigma \nabla_\nu \sigma + \frac{1}{6} R) \phi^2 + \frac{1}{2} \lambda \sigma \phi^4 \} + S_m[\psi, \bar{g}_{\mu\nu}] , \quad (4)$$

in which $\bar{g}_{\mu\nu} = e^\sigma g_{\mu\nu}$ and $S_m[\psi, \bar{g}_{\mu\nu}]$ stands for the matter action which contains some matter field variables, collectively denoted by ψ , with anomalous gravitational coupling. The new scalar field σ describes the distinction between the two conformal frames of the metric describing the gravitational and the matter part of the action (4). The presence of a kinetic term for σ implies that σ is considered as a dynamical field. The action (4) contains also an interaction term $\sim \lambda \sigma \phi^4$ in which λ is a dimensionless coupling constant. The particular form of this interaction term and the kinetic term of σ is suggested by the fact that these terms respect the conformal symmetry of the gravitational action (1)[‡]. Thus the total action (4) is distinguished by the fact that a conceivable (conformal) symmetry breaking may be linked to the physical characteristics of the matter at hand.

At this point it is necessary to clarify the notion of the anomalous gravitational coupling introduced by the action (4) with respect to other existing publications devoted to this issue. Usually one uses conformal transformations of the metric to transfer the action of a scalar-tensor theory into different inequivalent representations, for a review see [6]. This is done, for example, in Brans-Dicke theory to represent the corresponding action in the so called Einstein frame in which a constant strength of the gravitational coupling can be established. In this approach there is no a priori distinction between the conformal frame of the metric used in the gravitational part and that used in the matter part of the original action. The anomalous gravitational coupling emerges in all these cases because the applied conformal transformation generates an exponential factor before the metric [7] in the matter action or occasionally before the matter part of the Lagrangian [6]. The notion of the anomalous coupling introduced by the action (4) is however conceptually different with respect to two points. Firstly, it allows us to have an a priori distinction between the two conformal frames of the metric used in the gravitational part and in the matter part of the original action. Secondly, the action is constructed in such a way that the gravitational part of the action exhibits the conformal symmetry so that the physical characteristics of the matter may be considered as responsible for a conceivable conformal symmetry breaking.[§] This last point has important implications regarding a new symmetry breaking phase if the matter actually

[‡]Note that σ as a dimensionless function does not transform under a conformal transformation.

[§]Note that the class of scalar tensor theories which is mostly discussed in the context of the anomalous

breaks the conformal invariance. In such cases the anomalous gravitational coupling of matter can be related to an anomalous phase of the symmetry breaking. This anomalous phase is often ignored in the consideration of conformal symmetry breaking [8] which is usually based on the normal coupling (the normal phase). We shall deal with this issue later in connection with the symmetry breaking effect due to the gravitational coupling of the large-scale matter.

Varying S' with respect to $g^{\mu\nu}$, ϕ and σ yields, respectively,

$$G_{\mu\nu} - \frac{3}{2}\lambda\sigma\phi^2 g_{\mu\nu} = 6\phi^{-2}(e^\sigma \Sigma_{\mu\nu} + \tau_{\mu\nu}) + 6t_{\mu\nu} , \quad (5)$$

$$\square\phi - \frac{1}{6}R\phi - \lambda\sigma\phi^3 - \phi\nabla_\alpha\sigma\nabla^\alpha\sigma = 0 , \quad (6)$$

$$\nabla_\mu J^\mu = \frac{1}{2}\sqrt{-g}\left(\frac{1}{2}\lambda\phi^4 + e^\sigma g^{\mu\nu}\Sigma_{\mu\nu}\right) , \quad (7)$$

where

$$\Sigma_{\mu\nu} = \frac{-2}{\sqrt{\bar{g}}}\frac{\delta}{\delta\bar{g}^{\mu\nu}}S_m[\psi, \bar{g}_{\mu\nu}] , \quad (8)$$

$$J^\mu = \sqrt{-g}\phi^2 g^{\mu\nu}\nabla_\nu\sigma , \quad (9)$$

and

$$\tau_{\mu\nu} = -(\nabla_\mu\phi\nabla_\nu\phi - \frac{1}{2}g_{\mu\nu}\nabla_\alpha\phi\nabla^\alpha\phi) - \frac{1}{6}(g_{\mu\nu}\square - \nabla_\mu\nabla_\nu)\phi^2 , \quad (10)$$

$$t_{\mu\nu} = -\nabla_\mu\sigma\nabla_\nu\sigma + \frac{1}{2}g_{\mu\nu}\nabla_\alpha\sigma\nabla^\alpha\sigma . \quad (11)$$

Here $\square \equiv g^{\mu\nu}\nabla_\mu\nabla_\nu$ and $G_{\mu\nu}$ is the Einstein tensor of the metric $g_{\mu\nu}$. In the equation (5) we consider $\Sigma_{\mu\nu}$ as the matter stress tensor. Note that this tensor appears in (5) with a conformal weight e^σ . This is a consequence of the anomalous coupling which forces us to define $\Sigma_{\mu\nu}$ in terms of functional differentiation with respect to $\bar{g}_{\mu\nu}$ [¶].

Taking the four-divergence of (5) and combining the result with (6) and (7) leads to

$$\nabla^\mu(e^\sigma\Sigma_{\mu\nu}) = \frac{1}{2}\nabla_\nu\sigma e^\sigma g^{\mu\nu}\Sigma_{\mu\nu} , \quad (12)$$

which indicates that $\Sigma_{\mu\nu}$ can not be conserved due to the presence of σ . We also note that in the absence of matter the trace of (5) would lead directly to the equation (6) and, therefore, the latter would not contain any new information. This is a direct consequence of conformal invariance of the vacuum sector of (4). In the presence of matter, on the other hand, we obtain the following condition

$$e^\sigma g^{\mu\nu}\Sigma_{\mu\nu} = 0 , \quad (13)$$

by comparing the trace of (5) with the equation (6). This is a dynamical consistency relation imposed on the behavior of matter systems in this model. It means that only the gravitational

coupling breaks the conformal symmetry already in the vacuum sector, i.e., at the level of the gravitational part of the original action.

[¶]In the present context, there may be an ambiguity in the definition of the conformal weight of the matter stress tensor when matter systems anomalously couple to gravity. This ambiguity does not affect the basic result of the present work. The exploration of the issue is however deserved to be investigated elsewhere.

coupling of traceless matter systems is allowed to be described by (4). To describe the gravitational coupling of matter systems with nonvanishing trace of the corresponding stress-tensor one may add a term such as $\frac{1}{2} \int d^4x \sqrt{-g} \mu^2 \phi^2$ to the total action in (4) with μ being a constant mass parameter. In this case the analog of the equations (5) and (6) are

$$G_{\mu\nu} - 3\mu^2 g_{\mu\nu} - \frac{3}{2} \lambda \sigma \phi^2 g_{\mu\nu} = 6\phi^{-2} (e^\sigma \Sigma_{\mu\nu} + \tau_{\mu\nu}) + 6t_{\mu\nu} , \quad (14)$$

$$\square\phi - \frac{1}{6} R\phi - \lambda\sigma\phi^3 - \mu^2\phi - \phi\nabla_\alpha\sigma\nabla^\alpha\sigma = 0 , \quad (15)$$

and the equation (7) remains unchanged. The relation (13) takes then the form

$$e^\sigma g^{\mu\nu} \Sigma_{\mu\nu} = -\mu^2 \phi^2 . \quad (16)$$

This ensures the consideration of the gravitational coupling of matter systems with nonvanishing trace of the corresponding stress-tensor to be dynamically consistent.

It should however be noted that the appearance of a constant dimensional parameter such as μ clearly breaks the conformal symmetry. We are primarily interested in the investigation of a cosmological symmetry breaking emerging from the gravitational coupling of the large-scale distribution of matter in the universe. In this case the length scale μ^{-1} should be related to the typical size of the universe R_0 , namely $\mu^{-1} \sim R_0$. We first consider the normal phase of the symmetry breaking which is taken to be based on the normal coupling of the large-scale distribution of matter. This demands a vanishing average value for σ at large scales, namely $\bar{\sigma} = 0$. In this case it is possible to obtain an estimation for the constant background average value of ϕ which provides the strength of the gravitational coupling. In fact, the trace of $\Sigma_{\mu\nu}$ can be measured in terms of the average density of the large-scale distribution of matter, i.e., $\Sigma_\mu^\mu \sim -M/R_0^3$ which M denotes the mass of the universe. If one uses the empirical fact that the radius of the universe coincides with its schwarzschild radius $2GM$, one then gets an estimation of the constant background value of ϕ

$$\phi^{-2} \sim G \sim m_p^{-2} , \quad (17)$$

where G and m_p are the gravitational constant and the Planck mass, respectively. In this case, the gravitational equation (14) reduces to the Einstein field equations with a small cosmological constant. It is important to note that the use of $\lambda\sigma\phi^4$, instead of the usual $\lambda\phi^4$ term, allows to avoid the cosmological constant problem in a cosmological context.

Thus it is possible to use a constant configuration of ϕ to measure a constant strength of gravitational coupling of the large scale distribution of matter in the normal coupling. It should be emphasized that this feature seems to be characteristic only to the normal phase of the symmetry breaking which is based on the normal coupling. In the case of anomalous coupling the conformal symmetry breaking can not be used to achieve a precise estimation of the energy density of $\Sigma_{\mu\nu}$. This means that we may not obtain any knowledge of the configuration of the scalar field ϕ as a measure for the strength of the anomalous gravitational coupling.

We proceed now to the consideration of the anomalous phase of the symmetry breaking which is taken to be based on the anomalous coupling of the large-scale distribution of matter. The corresponding symmetry breaking is characterized by the relation (16) in which the

length scale μ^{-1} again measures the typical size of the universe. But (16) implies that the local variations of σ lead to a variable configuration of ϕ . As a consequence, the strength of the anomalous gravitational coupling of the large scale distribution of matter should not be considered as constant. This feature can be used to derive a particle concept in the anomalous phase of the symmetry breaking. From (15) we obtain

$$g^{\mu\nu}\nabla_\mu\sigma\nabla_\nu\sigma = -m^2 - \frac{1}{6}R + \frac{\square\phi}{\phi}, \quad (18)$$

which may be interpreted as a generalized Hamilton-Jacobi equation of a relativistic particle with a variable mass $m = (\lambda\sigma\phi^2 + \mu^2)^{1/2}$. The major characteristic of (18) lies in the term $\frac{\square\phi}{\phi}$. It is similar to the term related to the so-called quantum potential in the context of causal interpretation of relativistic quantum mechanics [2]. In present context, this term emerges because the anomalous gravitational coupling of large-scale matter systems does not provide us a constant strength for the gravitational coupling.

In order for the particle interpretation of (18) to be valid two general characteristics should be present. Firstly the particle trajectories must be timelike. Secondly the variable mass scale m must remain constant along the particle trajectories. To establish the first characteristic one may impose the condition $\nabla_\alpha\sigma\nabla^\alpha\sigma < 0$ to ensure that the particles move along timelike trajectories. Looking at (11) we see that this acts as an energy condition on the stress tensor $t_{\mu\nu}$. It selects, namely, those configurations of σ for which the trace of $t_{\mu\nu}$ is negative. For the stress tensors of matter systems this assumption is dynamically consistent, as one may infer from (16). Thus the restriction to timelike trajectories means that the stress tensor $t_{\mu\nu}$ which appears on the right hand side of (14) follows the general behavior of dynamically allowed matter systems with respect to the sign of the trace. The establishment of the second characteristic needs careful considerations. The constant mass scale of a particle seems to be a feature which is valid if a particle could ideally be considered as a quasi-isolated object. To perform the corresponding limiting procedure we propose to consider the scaling limit $\mu \rightarrow 0$, i.e., the idealized limit of infinite size of the universe. In this limit the conformal invariance is restored. Taking this limit as reasonable we may then select a conformal frame (a particle frame) by the condition

$$\nabla_\alpha\sigma\nabla^\alpha m = 0, \quad (19)$$

which ensures that the mass scale m remains constant along the particle trajectories^{||}.

Finally we turn to the general particle concept described by (18) and remark that the motion of the particles can be regarded as influenced by a guidance wave, or a pilot wave,

^{||}The masses of the particles contribute to vacuum energy density and appear as a large cosmological constant in the gravitational equation (14). Thus the question arises whether this would lead to a cosmological constant problem [9]. This issue is difficult to be resolved, but we make here only a modest remark. In the present case this problem arises only in the anomalous phase of symmetry breaking. In fact the mass of the particles is generated by a nonvanishing value of σ which in our presentation is characteristic to the anomalous coupling of the large-scale matter. However the considerations in cosmology are offer based on the normal coupling of the large-scale matter. The vanishing value of σ ensures in this case that no cosmological constant problem arises. One can probably say that the cosmological constant problem arises if one makes no distinction between the nature of gravitational coupling of matter in cosmology and particle physics [10]. The issue is however too difficult to be resolved within the scope of this paper.

defined by

$$\psi = \phi e^{i\sigma} . \quad (20)$$

Using (9) and (18), one can easily show that ψ satisfies the wave equation

$$(\square - m^2 - \frac{1}{6}R)\psi = \frac{i\phi^{-2}}{\sqrt{-g}} \nabla_\mu J^\mu \psi . \quad (21)$$

As a consequence of (7) and in the scaling limit $\mu \rightarrow 0$, this reduces to

$$(\square - m^2 - \frac{1}{6}R)\psi = i\frac{m^2}{4\sigma}\psi . \quad (22)$$

The right hand side of this equation indicates that there is an imaginary contribution to the squared mass of the particles. However one expects that such a contribution to be vanishingly small since in order to achieve a particle interpretation, there should exist a precise distinction between the two conformal frames of the metric describing the gravitational and the matter part of the action (4). Thus one may require that σ takes sufficiently large values. In this case the right hand side of the equation (22) drops out and this equation takes the form of the usual field equation in a gravitational background.

We summarize our observations as follows: A breakdown of conformal invariance of the gravitational model (4) in the anomalous phase provides a theoretical framework by which one may derive a particle concept. This derivation entails the existence of a dynamical field which describes the anomalous coupling of the large-scale matter, namely the scalar function σ . This scalar function makes it possible to have a distinction between the conformal frame of the metric used in the description of the large-scale matter system and that used in the description of the geometry of spacetime. In the absence of this distinction (the normal phase of symmetry breaking) the theory describes just the usual gravitational coupling of large-scale distribution of matter. In the presence of this distinction (the anomalous phase of symmetry breaking), on the other hand, the theory can be used to model local particle systems. In this case it is essentially the local variation of the scalar function σ that leads one to a particle interpretation in which σ plays the role of the Hamilton-Jacobi function. This demonstrates the nontrivial role of the absolute value of the phase of a wave function in the gravitational context. The local variation of σ has another important implication in that it predicts a variable strength of the gravitational coupling of the large-scale matter. This leads to the appearance of a corresponding term in the Hamilton-Jacobi equation which is similar to the quantum potential term used in the causal interpretation of quantum mechanics. The whole derivation of the particle concept suggests that the anomalous phase of the symmetry breaking due to the anomalous gravitational coupling of the large-scale matter may have a vital role to play in the geometric interpretation of causal framework of quantum mechanics.

References

- [1] J. D. Bekenstein, A. Meisels, Phys. Rev. D **22**, 1313 (1980)
- [2] D. Bohm, J. Hiley, The Undivided Universe, (Routledge 1993)
P. R. Holland, The Quantum Theory of Motion, (Cambridge University Press 1993)
- [3] W. R. Wood and G. Papini, Phys. Rev. D **45**, 3617 (1992)
- [4] W. R. Wood and G. Papini, Found. Phys. Lett. **6**, 207 (1993)
- [5] H. Salehi, H. Sepangi, F. Darabi, Found. Phys. Lett. **13**, 297 (2000)
- [6] V. Faraoni, E. Gunzig and P. Nardone, Funds. Cosmic Phys. **20**, 121 (1999)
- [7] T. Damour, G.W. Gibbons and C. Gundlach, Phys. Rev. Lett. **64**, 123 (1990)
- [8] S. Deser, Ann. Phys. **59**, 248 (1970)
- [9] S. Weinberg, Rev. Mod. Phys. **61**, 1 (1989)
- [10] Y. Bisabr, H. Salehi, Class. Quantum Grav. **19**, 2369 (2002)